arXiv:1103.3588v1 [math.CO] 18 Mar 2011

Characterization of n-Vertex Graphs with Metric Dimension $\mathbf{n-3}$

MOHSEN JANNESARI and BEHNAZ OMOOMI
Department of Mathematical Sciences
Isfahan University of Technology
84156-83111, Isfahan, Iran

January 15, 2013

Abstract

For an ordered set $W = \{w_1, w_2, \ldots, w_k\}$ of vertices and a vertex v in a connected graph G, the ordered k-vector $r(v|W) := (d(v, w_1), d(v, w_2), \ldots, d(v, w_k))$ is called the (metric) representation of v with respect to W, where $d(x, y)$ is the distance between the vertices x and y. The set W is called a resolving set for G if distinct vertices of G have distinct representations with respect to W. The minimum cardinality of a resolving set for G is its metric dimension. In this paper, we characterize all graphs of order n with metric dimension $n - 3$.

Keywords: Resolving set, Basis, Metric dimension, Locating set.

1 Introduction

Throughout this paper $G = (V, E)$ is a finite, simple, and connected graph of order $n(G)$. The distance between two vertices u and v, denoted by $d_G(u, v)$, is the length of a shortest path between u and v in G. We write it simply $d(u, v)$ when no confusion can arise. Also, the diameter of G, $max_{\{u,v\}\subseteq V(G)} d(u, v)$, is denoted by $diam(G)$. The vertices of a connected graph can be represented by different ways, for example, the vectors which theirs components are the distances between the vertex and the vertices in a given subset of vertices.

For an ordered set $W = \{w_1, w_2, \ldots, w_k\} \subseteq V(G)$ and a vertex v of G, the k-vector

$$r(v|W) := (d(v, w_1), d(v, w_2), \ldots, d(v, w_k))$$

is called the (metric) *representation* of v with respect to W. The set W is called a *resolving set* (locating set) for G if distinct vertices have different representations. In this case we say the set W *resolves* G. Elements in a resolving set are called *landmarks*. A resolving set W for G with minimum cardinality is called a *basis* of G, and its cardinality is the *metric dimension* of G, denoted by $\beta(G)$.

The concept of (metric) representation is introduced by Slater [11] (see [8]). For more results in this concept see [1, 2, 4, 5, 7, 13].

It is obvious that for every graph G of order n, $1 \leq \beta(G) \leq n-1$. Yushmanov [14] improved this bound to $\beta(G) \leq n - diam(G)$. Khuller et al. [10] and Chartrand et al. [6] independently proved that $\beta(G) = 1$ if and only if G is a path. Also, all graphs with metric dimension two are characterized by Sudhakara and Hemanth Kumar [12]. Chartrand et al. [6] proved that the only graph of order n, $n \geq 2$, with metric dimension $n-1$ is the complete graph K_n. They also provided a characterization of graphs of order n and metric dimension $n-2$. In [3] the problem of characterization of all graphs of order n with metric dimension $n-3$ is proposed. In this paper, we answer to this question and provide a characterization of graphs with metric dimension $n-3$. First in next section, we present some definitions and known results which are necessary to prove our main theorem.

2 Preliminaries

In this section, we present some definitions and known results which are necessary to prove our main theorem. The notations $u \sim v$ and $u \nsim v$ denote the adjacency and none-adjacency between vertices u and v, respectively. An edge with end vertices u and v is denoted by uv. A path of order n, P_n, and a cycle of order n, C_n, are denoted by $(v_1, v_2, \ldots, v_n)$ and $(v_1, v_2, \ldots, v_n, v_1)$, respectively.

We say an ordered set W *resolves* a set T of vertices in G, if the representations of vertices in T are distinct with respect to W. When $W = \{x\}$, we say that the vertex x resolves T. To see that whether a given set W is a resolving set for G, it is sufficient to look at the representations of vertices in $V(G)\backslash W$, because $w \in W$ is the only vertex of G for which $d(w,w) = 0$.

In [6] all graphs of order n with metric dimension $n-2$ are characterized as follows.

Theorem A. [6] *Let G be a graph of order $n \geq 4$. Then $\beta(G) = n-2$ if and only if $G = K_{s,t}$ $(s,t \geq 1)$, $G = K_s \vee \overline{K}_t$ $(s \geq 1, t \geq 2)$, or $G = K_s \vee (K_t \cup K_1)$ $(s,t \geq 1)$.*

For each vertex $v \in V(G)$, let $\Gamma_i(v) := \{u \in V(G) \mid d(u,v) = i\}$. Two distinct vertices u, v are *twins* if $\Gamma_1(v)\backslash\{u\} = \Gamma_1(u)\backslash\{v\}$. It is called that $u \equiv v$ if and only if $u = v$ or u, v are twins. In [9], it is proved that the relation $\equiv$ is an equivalent relation. The equivalence class of the vertex v is denoted by v^*. The *twin graph* of G, denoted by G^*, is the graph with vertex set $V(G^*) := \{v^* \mid v \in V(G)\}$, where $u^*v^* \in E(G^*)$ if and only if $uv \in E(G)$. It is easy to prove that u, v are adjacent in G if and only if all vertices of u^* are adjacent to all vertices of v^*, hence the definition of G^* is well defined. For each subset $S \subseteq V(G)$, let S^* denote the set $\{v^* \in V(G^*) \mid v^* \subseteq S\}$. Also, by $\Gamma_i^*(v^*)$, we mean the set $\{u^* \in V(G^*) \mid d_{G^*}(u^*, v^*) = i\}$. For each $v \in v^*$, it is immediate that $\Gamma_i^*(v^*) = \Gamma_i^*(v)$ if $i \neq 0$. Furthermore, we define

$$R_1(v) := \{x \in \Gamma_1(v) \mid \exists y \in \Gamma_2(v) : x \sim y\},$$

$$R_2(v) := \Gamma_1(v)\backslash R_1(v),$$

$$R_1^*(v^*) := \{x^* \in \Gamma_1^*(v^*) \mid \exists y^* \in \Gamma_2^*(v^*) : x^* \sim y^*\},$$

$R_2^*(v^*) := \Gamma_1^*(v^*) \backslash R_1^*(v^*)$.

We say a set S of vertices is *homogeneous* if the induced subgraph by S in G, $G[S]$, is a complete or an empty subgraph of G. In this terminology, it is proved in [9] that each vertex of G^* is a homogeneous subset of $V(G)$.

Observation 1. [9] *If the vertices u, v are twins in a graph G and S resolves G, then u or v is in S. Moreover, if $u \in S$ and $v \notin S$, Then $(S \setminus \{u\}) \cup \{v\}$ also resolves G.*

Proposition 1. [9] *Let $G \neq K_1$ be a graph. Then $diam(G^*) \leq diam(G)$. Moreover, if $u, v \in V(G)$ are not twin vertices of G, then $d_{G^*}(u^*, v^*) = d_G(u, v)$.*

As in [9], we say that $v^* \in V(G^*)$ is of type:

- (1) if $|v^*| = 1$,
- (K) if $G[v^*] \cong K_r$ and $r \geq 2$,
- (N) if $G[v^*] \cong \overline{K_r}$ and $r \geq 2$.

A vertex of G^* is of type (1K) if it is of type (1) or (K). A vertex is of type (1N) if it is of type (1) or (N). A vertex is of type (KN) if it is of type (K) or (N). We denote by $\alpha(G^*)$ the number of vertices of G^* of type (K) or (N). It is obvious that G is uniquely determined by G^*, and the type and cardinality of each vertex of G^*.

Hernando et al. [9] characterized all graphs of order n, diameter d and metric dimension $n-d$ by the following theorem.

Theorem B. [9] *Let G be a graph of order n and diameter $d \geq 3$. Let G^* be the twin graph of G. Then $\beta(G) = n - d$ if and only if G^* is one of the following graphs:*

1. *$G^* \cong P_{d+1}$ and one of the following cases holds*
   (a) $\alpha(G^*) \leq 1$;
   (b) $\alpha(G^*) = 2$, the two vertices of G^* not of type (1) are adjacent, and if one is a leaf of type (K), then the other is also of type (K);
   (c) $\alpha(G^*) = 2$, the two vertices of G^* not of type (1) are at distance 2 and both are of type (N); or
   (d) $\alpha(G^*) = 3$ and there is a vertex of type (KN) adjacent to two vertices of type (N).
2. *$G^* \cong P_{d+1,k}$ (the path $(u_0^*, u_1^*, \ldots, u_d^*)$ with one extra vertex adjacent to u_{k-1}^*) for some integer $k \in [3, d-1]$, the degree-3 vertex u_{k-1}^* of G^* is of any type, each neighbor of u_{k-1}^* is of type (1N), and every other vertex is of type (1).*
3. *$G^* \cong P'_{d+1,k}$ (the path $(u_0^*, u_1^*, \ldots, u_d^*)$ with one extra vertex adjacent to u_{k-1}^* and u_k^*) for some integer $k \in [2, d-1]$, the three vertices in the cycle are of type (1K), and every other vertex is of type (1).*

To prove our main theorem, we need the following propositions.

Proposition 2. *Let H be a subgraph of a graph G. If $\beta(H) = n(H) - t$ and $d_H(u,v) = d_G(u,v)$ for all pairs of vertices in H, then $\beta(G) \leq n(G) - t$.*

Proof. Let W be a basis of H. Since $d_H(u,v) = d_G(u,v)$ for all pairs of vertices in H, W as a subset of $V(G)$ resolves all vertices of H. Hence, $T = W \cup (V(G) \backslash V(H))$ is a resolving set for G where, $|T| = n(G) - t$. Therefore, the metric dimension of G is at most $n(G) - t$. ■

Corollary 1. *If H is an induced subgraph of a graph G, where $diam(H) = 2$, and $\beta(H) = n(H) - t$ for some positive integer t, then $\beta(G) \leq n(G) - t$.*

Corollary 2. *Let H be an induced subgraph of a graph G, and G be an induced subgraph of a graph R, where $diam(H) = diam(G) = 2$, $\beta(R) = n(R) - t$, and $\beta(H) = n(H) - t$ for some positive integer t. Then $\beta(G) = n(G) - t$.*

Proof. Let $\beta(G) = n(G) - s$, for some positive integer s. By Corollary 1, we have $n(G) - s = \beta(G) \leq n(G) - t$ and $n(R) - t = \beta(R) \leq n(R) - s$. Therefore, $s = t$. ■

Proposition 3. *Let G be a graph and G^* be the twin graph of G. If $\beta(G^*) = n(G^*) - t$ for some positive integer t, then $\beta(G) \leq n(G) - t$.*

Proof. Let S^* be a basis of G^* and $T^* = V(G^*) \backslash S^*$. We choose a vertex v from each $v^* \in T^*$ and let $T = \{v \mid v^* \in T^*\}$. Since S^* is a basis of G^*, for each pair of vertices $u^*, v^* \in T^*$, there exist a vertex $x^* \in S^*$ such that $d_{G^*}(x^*, u^*) \neq d_{G^*}(x^*, v^*)$. Note that neither u nor v is twin with x, for each $x \in x^*$. Therefore, by Proposition 1, we have $d_{G^*}(x^*, u^*) = d_G(x, u)$ and $d_{G^*}(x^*, v^*) = d_G(x, v)$. Hence, $d_G(x, u) \neq d_G(x, v)$, which implies that the set $S = \bigcup_{v^* \in S^*} v^*$ resolves T. Therefore, $V(G) \backslash T$ is a resolving set for G of cardinality $n(G) - t$, thus $\beta(G) \leq n(G) - t$. ■

Proposition 4. *Let G be a graph and G^* be the twin graph of G. Then $\beta(G) \geq n(G) - n(G^*)$.*

Proof. If S resolves G, then Observation 1 shows that S contains at least $|v^*| - 1$ vertices from each $v^* \in V(G^*)$. Hence, $\beta(G) \geq n(G) - n(G^*)$. ■

3 Main Results

Let G be a connected graph of order n and metric dimension $n - 3$. Since $n - 3 = \beta(G) \leq n - diam(G)$, it follows that $diam(G) \leq 3$. If $diam(G) = 1$, then $G \cong K_n$, contrary to $\beta(G) = n - 3$. If $diam(G) = 3$, then in Theorem B, let $d = 3$, which obtains a characterization of graphs G with $\beta(G) = n - 3$, where $diam(G) = 3$, (Note that in this case the interval $[3, 2]$ is empty, hence, Case 2 dose not occur). Therefore, it is enough to consider the case $diam(G) = 2$. The following theorem is our main result, which is a characterization of all graphs with metric dimension $n-3$ and diameter 2.

Theorem 1. *Let G be a graph of order n and diameter 2 and G^* be the twin graph of G. Then $\beta(G) = n-3$ if and only if G^* satisfies in one of the following structures:*

G_1*: $G^* \cong K_3$ and has at most one vertex of type (1K);*

G_2*: $G^* \cong P_3$ and one of the following cases holds:*
(a) The degree-2 vertex is of type (N) and one of the leaves is of type (K) and the other is of any type;
(b) One of the leaves is of type (K), the other is of type (KN) and the degree-2 vertex is of any type;

G_3*: G^* is the paw (a triangle with a pendant edge), and the degree-3 vertex is of any type, one of the degree-2 vertices is of type (N), the other is of type (1K), the leaf is of type (1N). Moreover, a degree-2 vertex of type (K) yields the leaf and the degree-3 vertex are not of type (N);*

G_4*: $G^* \cong C_5$, and each vertex is of type (1);*

G_5*: G^* is C_5 with a chord, and the adjacent degree-2 vertices are of type (1) and the other vertices are of type (1K);*

G_6*: G^* is a C_5 with two adjacent chords. The degree-4 vertex is of any type, the others are of type (1K). Furthermore, two non-adjacent vertices are not of type (K), and two adjacent vertices are not of different types (K) and (N);*

G_7*: G^* is a kite with a pendant edge adjacent to a degree-3 vertex, and the leaf is of type (1), the degree-4 and degree-3 vertices are of type (1K), one of the degree-2 vertices is of type (K) and the other is of type (1).*

G_8*: G^* is the kite, and one of the degree-2 vertices is of type (K) the other is of type (1), one of the degree-3 vertices is of type (N), and the other is of type (1K);*

G_9*: $G^* \cong C_4$, and two adjacent vertices are of type (K) and others are of type (1);*

G_{10}*: $G^* \cong C_4 \vee K_1$, and two degree-3 adjacent vertices are of type (K), degree-4 vertex is of type (1K), and others are of type (1).*

In Figure 1 the scheme of the above 10 structures are shown.

3.1 Proof of Necessity

Throughout this section, G is a graph of order n, diameter 2, metric dimension $n-3$, and G^* is the twin graph of G. Note that, Proposition 1 implies that $diam(G^*) \leq 2$. Through a sequence of lemmas and propositions, we show that G^* has one of the structures G_1 to G_{10}.



Figure 1: The twin graphs of graphs with diameter 2 and metric dimension $n-3$.

Proposition 5. *If $diam(G^*) = 1$, then G^* satisfies in structure G_1.*

Proof. Let u^*, v^* be two vertices of G^* of type (1K). Since G^* is a complete graph, every pair of vertices $u \in u^*$ and $v \in v^*$ of G are twins, thus $u^* = v^*$. Hence, G^* has at most one vertex of type (1K). If vertices v_1^*, v_2^*, v_3^*, and v_4^* of G^* (except possibly v_1^*) are of type (N), then we choose an arbitrary vertex $v_i \in v_i^*$, for each i, $1 \leq i \leq 4$, and $u_i \in v_i^* \backslash \{v_i\}$, for each i, $2 \leq i \leq 4$. Let $T = \{u_2, u_3, u_4\}$. It follows that

$$r(v_1|T) = (1,1,1), \quad r(v_2|T) = (2,1,1), \quad r(v_3|T) = (1,2,1), \quad r(v_3|T) = (1,1,2).$$

Hence, the set $V(G)\backslash\{v_1, v_2, v_3, v_4\}$ is a resolving set for G and $\beta(G) \leq n-4$. This contradicts our assumption, $\beta(G) = n-3$. Therefore, G^* has at most three vertices. Assume $G^* \cong K_t$ for some integer $t \in [1,3]$. If $t = 1$, then $G \cong K_n$ or $G \cong \overline{K}_n$. If $t = 2$, then $G \cong K_{r,s}$ or $G \cong K_r \vee \overline{K}_s$. Since $\beta(G) = n-3$, the above cases are impossible. Consequently $G^* \cong K_3$, which is the desired conclusion. ■

The remainder of this section will be devoted to the case $diam(G^*) = 2$. It is clear that in this case, there exists a vertex $v \in V(G)$ such that $\Gamma_2^*(v^*) \neq \emptyset$ and $\Gamma_2(v) \neq \emptyset$.

Lemma 1. *If $\Gamma_2^*(v^*) \neq \emptyset$ and $v \in v^*$, then $\bigcup_{u^* \in \Gamma_i^*(v^*)} u^* \subseteq \Gamma_i(v)$, where $i \in \{1,2\}$, $\bigcup_{u^* \in R_2^*(v)} u^* \subseteq R_2(v)$, and $\bigcup_{u^* \in R_1^*(v)} u^* = R_1(v)$. Moreover, if v^* is of type (1K), then $\bigcup_{u^* \in \Gamma_2^*(v^*)} u^* = \Gamma_2(v)$, $R_1^*(v^*) = R_1^*(v)$, and $R_2^*(v^*) = R_2^*(v)$.*

Proof. It is clear that a vertex in $\Gamma_1(v)$ is not twin with any vertex of $\Gamma_2(v)$. Therefore, all twins of vertices in $\Gamma_1(v)$ are in the set $\Gamma_1(v) \cup \{v\}$ and all twins of vertices in $\Gamma_2(v)$ are in $\Gamma_2(v) \cup \{v\}$. This gives $\Gamma_i(v)\backslash v^* = \bigcup_{u^* \in \Gamma_i^*(v^*)} u^*$, for $i \in \{1,2\}$. Hence, $\bigcup_{u^* \in \Gamma_i^*(v^*)} u^* \subseteq \Gamma_i(v)$, when $i \in \{1,2\}$.

Note that all twins of vertices in $R_1(v)$ are in $R_1(v)$, because each member of $R_1(v)$ is adjacent to v and has at least one neighbor in $\Gamma_2(v)$ while the vertices in $R_2(v) \cup \{v\}$ have not such neighbors. Thus $\bigcup_{u^* \in R_1^*(v)} u^* = R_1(v)$. By the same reason, all twins of vertices in $R_2(v)$ are in $R_2(v) \cup \{v\}$. Consequently, $\bigcup_{u^* \in R_2^*(v)} u^* = R_2(v) \backslash v^* \subseteq R_2(v)$.

Now let v^* is of type (1K). Therefore, v can only be twin with the vertices of $R_2(v)$. Hence, $\bigcup_{u^* \in \Gamma_2^*(v^*)} u^* = \Gamma_2(v)$. It is clear that $R_1^*(v^*) \subseteq R_1^*(v)$. If there exist a vertex $u^* \in R_1^*(v) \backslash R_1^*(v^*)$, then $u^* \in R_2^*(v^*)$, because $\Gamma_1^*(v^*) = R_1^*(v^*) \cup R_2^*(v^*)$. Since $v^* \subseteq R_2(v) \cup \{v\}$, all neighbors of each vertex $u \in u^*$ are in the set $\Gamma_1(v) \cup \{v\}$, this contradicts the fact that $u \in R_1(v)$. Therefore, $R_1^*(v^*) = R_1^*(v)$ and consequently, $R_2^*(v^*) = R_2^*(v)$. ■

Lemma 2. *For each v, where $\Gamma_2(v) \neq \emptyset$, at least one of the sets $\Gamma_1(v)$ and $\Gamma_2(v)$ is homogeneous.*

Proof. Let $\Gamma_2(v) \neq \emptyset$. On the contrary, suppose that both $\Gamma_1(v)$ and $\Gamma_2(v)$ are not homogeneous. Therefore, there exist vertices v_1, v_2, and v_3 in $\Gamma_1(v)$, and vertices u_1, u_2, and u_3 in $\Gamma_2(v)$ such that $v_1 \sim v_2, v_2 \nsim v_3$ and $u_1 \sim u_2$, $u_2 \nsim u_3$. If $W' = \{v, v_2, u_2\}$, then the representations of v_1, v_3, u_1 and u_3 with respect to W' are as follows

$$r(v_1|W') = (1,1,*), \quad r(v_3|W') = (1,2,*), \quad r(u_1|W') = (2,*,1), \quad r(u_3|W') = (2,*,2),$$

where $*$ is 1 or 2. These four representations are distinct, hence, $V(G) \backslash \{v_1, v_3, u_1, u_3\}$ is a resolving set for G. Thus $\beta(G) \leq n-4$, which is a contradiction. Therefore, at least one of the sets $\Gamma_1(v)$ or $\Gamma_2(v)$ is homogeneous. ■

By Lemma 2, to complete the proof of necessity, we need to consider the following two cases.

Case 1. There exist a vertex $v \in V(G)$ such that $\Gamma_2^*(v^*) \neq \emptyset$ and $\Gamma_1(v)$ is homogeneous.

By the assumption of Case 1, the following results are obtained.

Fact 1. $|R_1^*(v^*)| \leq 2$.

Proof. Since every vertex of $R_1^*(v^*)$ has a neighbor in $\Gamma_2^*(v^*)$ and $\Gamma_1(v)$ is homogeneous, for distinct vertices $x^*, y^* \in R_1^*(v^*)$, the sets $\Gamma_1^*(x^*) \bigcap \Gamma_2^*(v^*)$ and $\Gamma_1^*(y^*) \bigcap \Gamma_2^*(v^*)$ are distinct nonempty sets. Therefore, the set $\Gamma_2^*(v^*)$ resolves the vertices of $R_1^*(v^*)$ in G^*. Moreover, since every vertex of $R_1^*(v^*)$ has a neighbor in $\Gamma_2^*(v^*)$ and v^* has not such a neighbor, if we compute the representations of the vertices in $R_1^*(v^*) \cup \{v^*\}$ with respect to $\Gamma_2^*(v^*)$, then the representation of each vertex in $R_1^*(v^*)$ has a coordinate 1 while all coordinates of $r(v^*|\Gamma_2^*(v^*))$ are 2. Therefore, $\Gamma_2^*(v^*)$ resolves the set $R_1^*(v^*) \cup \{v^*\}$ consequently, $V(G^*) \backslash (R_1^*(v^*) \cup \{v^*\})$ is a resolving set for G^*. Thus $\beta(G^*) \leq n(G^*) - |R_1^*(v^*) \cup \{v^*\}|$. On the other hand, by Propositions 3, we have $\beta(G^*) \geq n(G^*) - 3$. Therefore, $|R_1^*(v^*) \cup \{v^*\}| \leq 3$, which yields $|R_1^*(v^*)| \leq 2$. ■

Lemma 3. *If $\Gamma_2(v)$ is not homogeneous, then $R_2(v)$ and $R_2^*(v^*)$ are empty sets.*

Proof. Note that $|R_1^*(v^*)| \geq 1$, otherwise $\Gamma_2^*(v^*) = \emptyset$. Since $\Gamma_2(v)$ is not homogeneous, there exist vertices i, j, and k in Γ_2 such that $i \sim j$ and $j \nsim k$. Now if $R_2(v) \neq \emptyset$, then let $r_1 \in R_1(v) \cap \Gamma_1(j)$, $r_2 \in R_2(v)$ and $W_0 = \{v, j\}$. Thus, $r(i|W_0) = (2,1)$, $r(k|W_0) = (2,2)$, $r(r_1|W_0) = (1,1)$ and $r(r_2|W_0) = (1,2)$, and so $\beta(G) \leq n-4$. This contradiction implies $R_2(v) = \emptyset$.

If $R_2^*(v^*) \neq \emptyset$, then $\bigcup_{u^* \in R_2^*(v^*)} u^* \nsubseteq R_2(v)$ and hence, by Lemma 1, v^* is of type (N). Therefore, there exist two adjacent vertices $a, b \in \bigcup_{u^* \in \Gamma_2^*(v^*)} u^*$, otherwise $\Gamma_2(v)$ is homogeneous. Since $diam(G^*) = 2$, there exist a vertex $r_1 \in \bigcup_{u^* \in R_1^*(v^*)} u^*$ such that $r_1 \sim a$. Now let $v_1, v_2 \in v^*$, $r_2 \in \bigcup_{u^* \in R_2^*(v^*)} u^*$, and $W = \{v_1, a\}$. Thus, we have

$$r(v_2|W) = (2,2), \quad r(r_1|W) = (1,1), \quad r(r_2|W) = (1,2), \quad r(b|W) = (2,1).$$

Therefore, $V(G) \backslash \{v_2, r_1, r_2, b\}$ is a resolving set for G, with cardinality $n-4$, which contradicts our assumption $\beta(G) = n-3$. Consequently $R_2^*(v^*) = \emptyset$. ■

Fact 2. $|\Gamma_2^*(v^*)| \leq 3$.

Proof. If $\Gamma_2(v)$ is homogeneous, then since every vertex of $\Gamma_2^*(v^*)$ has a neighbor in $R_1^*(v^*)$ and $\Gamma_2^*(v^*)$ is homogeneous, for distinct vertices $x^*, y^* \in \Gamma_2^*(v^*)$, the sets $\Gamma_1^*(x^*) \bigcap R_1^*(v^*)$ and $\Gamma_1^*(y^*) \bigcap R_1^*(v^*)$ are distinct nonempty sets. Therefore, for each pair of vertices $x^*, y^* \in \Gamma_2^*(v^*)$ there exist a vertex $r_1^* \in R_1^*(v^*)$ such that r_1^* resolves x^* and y^* in G^*. Hence, $R_1^*(v^*)$ resolves all vertices of the set $\Gamma_2^*(v^*)$. This implies that $V(G^*) \backslash \Gamma_2^*(v^*)$ is a resolving set for G^*, which yields $\beta(G^*) \leq n(G^*) - |\Gamma_2^*(v^*)|$. On the other hand, by Propositions 3, we have $\beta(G^*) \geq n(G^*) - 3$. Thus, $|\Gamma_2^*(v^*)| \leq 3$.

Now let $\Gamma_2(v)$ is not homogeneous. By Fact 1, $|R_1^*(v^*)| \leq 2$. Thus, we consider two cases, $|R_1^*(v^*)| = 1$ and $|R_1^*(v^*)| = 2$. In the case $|R_1^*(v^*)| = 1$, let $R_1^*(v^*) = \{r_1^*\}$, $r_1 \in r_1^*$, and for each $l \in \Gamma_2(v)$, $N_1(l) = \Gamma_1(l) \cap \Gamma_2(v)$, and $N_2(l) = \Gamma_2(l) \cap \Gamma_2(v)$. Since $\Gamma_2(v)$ is not homogeneous, there exists a vertex $x \in \Gamma_2(v)$ such that both $N_1(x)$ and $N_2(x)$ are nonempty sets. Let $a \in N_1(x)$ and $y \in N_2(x)$. Note that $\Gamma_2^*(v^*) = \{x^*\} \cup N_1^*(x) \cup N_2^*(x)$, and x resolves a and y. Hence, if $N_1(x)$ is not homogeneous, then there exist vertices $i, j, k \in N_1(x)$ such that $i \sim j$ and $j \nsim k$. Thus, $W' = \{v, x, j\}$ resolves the $\{i, k, y, r_1\}$, this contradiction yields $N_1(x)$ is homogeneous. By a similar reason $N_2(x)$ is also homogeneous. Note that all different neighbors of vertices in $N_1^*(x)$ are in the set $N_2^*(x)$, because $N_1(x)$ is homogeneous and its vertices share their neighbors in $\Gamma_1(v) \cup \{x\}$. Similarly every different neighbor of vertices in $N_2^*(x)$ are in $N_1^*(x)$, hence, $N_1^*(x)$ and $N_2^*(x)$ resolves each other. Now let $W_1 = N_2^*(x) \cup \{x^*\}$. If each vertex of $N_1^*(x)$ has a none-neighbor vertex in $N_2^*(x)$, then the representation of each vertex of $N_1^*(x)$ with respect to $N_2^*(x)$ has a coordinate 2, all coordinates of $r(r_1^*|N_2^*(x))$ is 1, and $r(v^*|N_2^*(x))$ is entirely 2. Consequently, W_1 resolves $N_1^*(x) \cup \{r_1^*, v^*\}$. Thus, $\beta(G) = n-3$ implies that $|N_1^*(x)| \leq 1$. Moreover, if there exist a vertex $a^* \in N_1^*(x)$ such that a^* is adjacent to all vertices of $N_2^*(x)$, then $N_1^*(x)$ has at most two vertices. Otherwise, there are two distinct vertices $b^*, c^* \in N_1^*(x)$ such that they are different from a^*, and $r(b^*|N_2^*(x))$ and $r(c^*|N_2^*(x))$ are not entirely 1, and so W_1 resolves four vertices a^*, b^*, c^*, and v^*, contrary to $\beta(G) = n-3$. Hence, $|N_1^*(x)| \leq 2$. Furthermore, $|N_1^*(x)| = 2$ yields that there exist a vertex $a^* \in N_1^*(x)$ such that a^* is adjacent to all vertices of $N_2^*(x)$. On a similar way $|N_2^*(x)| \leq 2$, moreover, $|N_2^*(x)| = 2$ only if there exist a vertex $y^* \in N_2^*(x)$ such that y^* is none-adjacent to all vertices of $N_1^*(x)$. Thus, at most one of the sets $N_1^*(x)$ and $N_2^*(x)$ can have two vertices, because it is impossible that there exist a pair of vertices $a^* \in N_1^*(x),\ y^* \in N_2^*(x)$ such that a^* is adjacent to all vertices of $N_2^*(x)$ and y^* is none-adjacent to all vertices of $N_1^*(x)$. Consequently $|\Gamma_2^*(v^*)| \leq 4$. We claim that $|\Gamma_2^*(v^*)| = 4$ is impossible.

If $|\Gamma_2^*(v^*)| = 4$, then one of the two blow cases can be happened.

1. $|N_1^*(x)| = 2$ and $|N_2^*(x)| = 1$. Let $N_1^*(x) = \{a^*, b^*\}$, $N_2^*(x) = \{y^*\}$, $y^* \sim a^*$, $v \in v^*, a \in a^*, b \in b^*, x \in x^*$ and $y \in y^*$. If $a^* \sim b^*$, then x^* and b^* are twins, see Figure 2(a). Since $x^* \sim b^*$, one of them is of type (N). Note that b^* is not of type (N), because $N_1^*(x)$ is homogeneous and $a^* \sim b^*$. Hence, b^* is of type (1K) and x^* is of type (N). Thus, the set $V(G)\backslash\{r_1, y, a, x\}$ is a resolving set for G of size $n-4$, which is impossible. Therefore, $a^* \nsim b^*$, this gives the set $V(G)\backslash\{v, r_1, x, a\}$ is a resolving set for G. Which is a contradiction.

2. $|N_1^*(x)| = 1$ and $|N_2^*(x)| = 2$. Let $N_1^*(x) = \{a^*\}$, $N_2^*(x) = \{y^*, z^*\}$, $z^* \sim a^*$, $v \in v^*, a \in a^*, x \in x^*, y \in y^*, z \in z^*$ and $S = \{x, y\}$. Thus, $r(z|S) = (2, *)$, $r(a|S) = (1, 2)$, $r(r_1|S) = (1, 1)$ and $r(v|S) = (2, 2)$, where $* = 1$ or 2. Note that $\beta(G) = n-3$ yields $* = 2$, see Figure 2(b). Therefore, x^* and z^* are twins. Since they are none-adjacent vertices, one of them is of type (K). Clearly z^* is not of type (K), otherwise, since $N_2^*(x)$ is homogeneous, we have $* = 1$, which is impossible. Consequently, x^* is of type (K), this implies that the set $(x^*\backslash\{x\}) \cup \{z, y\}$ resolves $\{v, r_1, a, x\}$. Thus, $\beta(G) \leq n-4$.

These contradictions yield, $|\Gamma_2^*(v^*)| \leq 3$, when $|R_1^*(v^*)| = 1$.

To complete the proof we need only to consider the case $|R_1^*(v^*)| = 2$. In this case, since all different neighbors of vertices in $R_1(v)$ are in $\Gamma_2(v)$, $|R_1^*(v^*)| = 2$ implies that there exist a vertex $a \in \Gamma_2(v)$ such that $\Gamma_2(a) \cap R_1(v) \neq \emptyset$. Let $T = \Gamma_2(v)\backslash\{a\}$. Since $\Gamma_2(v)$ is not homogeneous, it has at least three vertices, hence $|T| \geq 2$. If T is not homogeneous, then there exists a vertex $i \in T$ such that i resolves two vertices of T. Moreover, we know that a resolves two vertices of $R_1(v)$. Hence, $\{v, i, a\}$ resolves at least four vertices of G. Thus, we obtain a resolving set for G of size $n-4$, which is impossible. Therefore, T is homogeneous.

If a is adjacent to a vertex in T, then a is adjacent to all vertices of T, otherwise $\{a, v\}$ resolves four vertices. If a vertex $t \in T$ is not adjacent to some vertices of $R_1(v)$, then similar to above, it can be seen that $\Gamma_2(v)\backslash\{t\}$ is homogeneous and t is adjacent or none-adjacent to all vertices of $\Gamma_2(v)\backslash\{t\}$. This implies that $\Gamma_2(v)$ is homogeneous, which is a contradiction with the assumption. Hence, every vertex of T is adjacent to all vertices of $R_1(v)$. Therefore, all vertices of T are twins and form a vertex b^* in G^*. Thus, $\Gamma_2^*(v^*)$ consists of two vertices a^* and b^*, where a^* is of type (1) and b^* is of type (KN). ■

Proposition 6. *If $|R_1^*(v^*)| = 1$, then G^* satisfies one of the structures G_2, G_3, or G_7.*

Proof. Let $R_1^*(v^*) = \{r_1^*\}$. First let $\Gamma_2(v)$ is homogeneous. Therefore, every vertex of $\Gamma_2^*(v^*)$ is adjacent to r_1^* and all vertices of $\Gamma_2^*(v^*)$ are twins. Consequently, since $\Gamma_2(v)$ is homogeneous, all vertices of $\bigcup_{u^* \in \Gamma_2^*(v^*)} u^*$ are twins. This gives $|\Gamma_2^*(v^*)| = 1$. Now, If $R_2^*(v^*) = \emptyset$, then $G^* \cong P_3$ and $\alpha(G^*) \geq 2$, otherwise $\beta(G) = n-2$. It is easy to check that in both cases $\alpha(G^*) = 2$ and 3, at



Figure 2: Different cases of $N_1^*(x)$ and $N_2^*(x)$.

least one of the leaves is of type (K), otherwise $\beta(G) = n-2$. Let $G^* = (x_1^*, x_2^*, x_3^*)$ and x_1^* be of type (K). If x_2^* is of type (1K) and x_3^* is of type (1), then $\beta(G) = n-2$. This contradiction implies that, if x_3^* is of type (1), then x_2^* is of type (N). That implies G^* satisfies the structure G_2.

If $R_2^*(v^*) \neq \emptyset$, then all vertices of $R_2^*(v^*)$ have a neighbor in $R_1^*(v^*)$, otherwise $diam(G^*) \geq 3$. Since $\Gamma_1(v)$ is homogeneous, every vertex of $R_2^*(v^*)$ is adjacent to every vertex of $R_1^*(v^*)$. Hence, all vertices of $R_2^*(v^*)$ are twins. This implies that $\Gamma_1(v)$ is a clique, all vertices of $\bigcup_{u^* \in R_2^*(v^*)} u^*$ are twins and so $|R_2^*(v^*)| = 1$. In this case G^* is the paw and one of the degree-2 vertices is v^* and the other belongs to $\Gamma_1^*(v^*)$. Therefore, the structure of G^* is as Figure 3(a). Hence, x^* and v^* are adjacent twins. Therefore, one of them is of type (N), otherwise $x^* = v^*$, which contradicts $R_2^*(v^*) \neq \emptyset$. Since $\Gamma_1(v)$ is a clique, x^*, y^* are of type (1K). Thus, v^* is of type (N). If z^* is of type (K), then we choose arbitrary fixed vertices $x \in x^*, v_1, v_2 \in v^*, y \in y^*$ and $z_1, z_2 \in z^*$. Hence,

$$r(v_1|\{v_2, z_2\}) = (2,2),\ r(x|\{v_2, z_2\}) = (1,2),\ r(y|\{v_2, z_2\}) = (1,1),\ r(z_1|\{v_2, z_2\}) = (2,1).$$

Thus, the set $V(G)\backslash\{x, y, z_1, v_1\}$ is a resolving set for G, and $\beta(G) \leq n-4$. Consequently, z^* is of type (1N). Similarly if x^* is of type (K) and z^* is of type (N), then $V(G)\backslash\{x, y, z_1, v_1\}$ is a resolving set for G that is a contradiction. Hence, G^* satisfies the structure G_3.

Now let $\Gamma_2(v)$ is not homogeneous. By the same notation as the proof of Fact 2, we can see that the vertices of $N_1(x)$ can only be twins with each other and x, also the vertices of $N_2(x)$ can only be twins with each other, x, and v. By Lemma 3, we have $R_2^*(v^*) = \emptyset$. Hence, if $|\Gamma_2^*(v^*)| = 1$, then $G^* \cong P_3$, all vertices of $N_2(x)$ are twins with v, and all vertices of $N_1(x)$ are twins with x. Thus, x^* is of type (K), v^* is of type (N), and r_1^* is of any type. In this case G^* satisfies the structure G_2.

When $|\Gamma_2^*(v^*)| = 2$, there exist three cases.

1. The vertex x and all vertices of $N_1(x)$ are twins and there exist some vertices in $N_2(x)$ which are not twins with v and they are twins by themselves. Thus, the vertices of $N_2(x)\backslash v^*$ form exactly one vertex, y^* in G^*. Hence, x^* is of type (K) and $x^* \nsim y^*$. Therefore, y^* and v^* are twins. Since $v^* \nsim y^*$, one of them is of type (K). Note that, if v^* is of type (K), then there exist some vertex $u \in V(G)$ which is adjacent to all vertices of $\Gamma_1(v) \cup \{v\}$ and is not adjacent to any vertex of $\Gamma_2(v)$. Hence, $u \in R_2(v)$, which is impossible, because by Lemma 3, $R_2(v) = \emptyset$. Thus, y^* is of

type (K). Let $r_1 \in r_1^*$, $y \in y^*$, $x \in x^*$ and $v \in v^*$. Then $(x^* \backslash \{x\}) \cup (y^* \backslash \{y\})$ resolves $\{v, r_1, x, y\}$, this contradicts our assumption $\beta(G) = n-3$. Consequently, this case does not happen.

2. There exist some vertices of $N_2(x)$ which are twins with x and the rest are twins with v, and all vertices of $N_1(x)$ are twins. Therefore, the vertices of $N_1(x)$ create a vertex a^* in G^*, $a^* \sim x^*$, and x^* is of type (N). Hence, G^* is the paw, with the leaf v^*, the degree-3 vertex r_1^*, and degree-2 vertices a^* and x^*. If a^* is of type (N), then $V(G)\backslash\{v, r_1, x, a\}$ is a resolving set for G, where $r_1 \in r_1^*, v \in v^*, x \in x^*$ and $a \in a^*$. This contradiction yields a^* is of type (1K). Because $R_2(v) = \emptyset$, v^* is not of type (K). Therefore, v^* is of type (1N). By a similar method as before, we deduce that, if a^* is of type (K), then v^* and r_1^* can not be of type (N). Thus, G^* satisfies in structure G_3.

3. All vertices of $N_2(x)$ are twins with v, and there exist some vertices in $N_1(x)$ which are not twins with x. Hence, the vertices of $N_1(x)\backslash x^*$ form a unique vertex a^* in G^*, $a^* \sim x^*$, and consequently G^* is the paw. Note that v^* is the leaf and its type is (N), the vertex r_1^* has degree 3, and x^*, a^* are degree-2 vertices. Since x^* and a^* are adjacent twins, one of them is of type (N). Also, since all vertices of $N_2^*(x)$ are twins with v, x^* is of type (1K), and so a^* is of type (N). Therefore, G^* has the structure G_3.

Finally, if $|\Gamma_2^*(v^*)| = 3$, then we have three following cases.

1. Every vertex of $N_2(x)$ is twin with v or x, and $N_1^*(x)$ has two vertices a^* and b^*. In this case a^* and b^* are twins, hence $a^* = b^*$, because $N_1(x)$ is homogeneous. Thus, $|\Gamma_2^*(v^*)| \leq 2$. Therefore this is not the case.

2. Every vertex of $N_1(x)$ is twin with x, and $N_2^*(x)$ has two vertices y^* and z^*. Hence, y^* and z^* are twins. Since $N_2(x)$ is homogeneous, $y^* = z^*$, which is a contradiction with $|\Gamma_2^*(v^*)| = 3$.

3. There exist some vertices in $N_1(x)$ which are not twins with x, also there exist some vertices in $N_2(x)$ which are neither twins with x nor v. In this case, each one of $N_1^*(x)$ and $N_2^*(x)$ has exactly one vertex a^* and y^*, respectively. If $a^* \nsim y^*$, then v^* and y^* are none-adjacent twins. Hence one of them is of type (K). Since $R_2(v) = \emptyset$, v^* is of type (1N), and so y^* is of type (K). Let $v \in v^*$, $y \in y^*$, $a \in a^*$, and $x \in x^*$. Then $V(G)\backslash\{y, x, r_1, v\}$ is a resolving set for G. This contradiction yields $a^* \sim y^*$, and in consequence, G^* is a kite with a pendant edge, adjacent to a degree-3 vertex. Thus, y^* and x^* are none-adjacent twins, hence one of them is of type (K). By symmetry, we can assume x^* is of type (K). Since $R_2(v) = \emptyset$, v^* is of type (1N). As we see before, v^*, a^*, and r_1^* are not of type (N) and y^* is not of type (KN). Therefore, G^* satisfies the structure G_7 and the proof is completed. ■

Heretofore we have considered the case $|R_1^*(v^*)| = 1$. Now we investigate the case $|R_1^*(v^*)| = 2$.

Proposition 7. *If* $|R_1^*(v^*)| = 2$ *and* $\Gamma_2(v)$ *is not homogeneous, then* G^* *satisfies the structure* G_7.

Proof. By the same notation as the proof of Fact 2, let $\Gamma_2^*(v^*) = \{a^*, b^*\}$, where a^* is of type (1) and b^* is of type (KN). Thus, if $a^* \sim b^*$, then b^* is of type (N), however $a^* \nsim b^*$ implies that b^* is of type (K), because $\Gamma_2(v)$ is not homogeneous. Let $R_1^*(v^*) = \{x^*, y^*\}$. Since $R_1(v)$ has a none-adjacent vertex to a, the vertex a^* has exactly one neighbor in $R_1^*(v^*)$. By the proof of



(a) (b)

Figure 3: $|\Gamma_2^*(v^*)| = 1$ and $|\Gamma_2^*(v^*)| = 2$.

Fact 2, $b^* \sim x^*$ and $b^* \sim y^*$, thus G^* is the 4-cycle, $C_4 = (v^*, x^*, b^*, y^*, v^*)$ with the pendant edge x^*a^* and possibly extra edges a^*b^* and x^*y^*, see Figure 3(b). Because $diam(G^*) = 2$, at least one of the edges a^*b^* and x^*y^* exists. If $a^* \sim b^*$, then b^* is of type (N). Let $v \in v^*$, $a \in a^*$, $b \in b^*$, $x \in x^*$, and $y \in y^*$. Consequently, the set $a^* \cup (b^* \backslash \{b\})$ resolves $\{v, x, b, y\}$, since b^* is of type (N). This contradiction yields $a^* \nsim b^*$, and so $x^* \sim y^*$, a^* is of type (1), and b^* is of type (K). Note that x^* and y^* are not of type (N), otherwise $\Gamma_1(v)$ is not homogeneous. Also, we can see easily that v^* is not of type (KN). In this case G^* satisfies the structure G_7. ■

Now, we need only to consider the case $|R_1^*(v^*)| = 2$ when $\Gamma_2(v)$ is homogeneous. In this case, if $|\Gamma_2^*(v^*)| = 1$, then all vertices of $R_1(v)$ are twins and consequently, $|R_1^*(v)| = 1$, which contradicts $|R_1^*(v^*)| = 2$. Therefore, $|\Gamma_2^*(v^*)| \geq 2$.

Lemma 4. *If* $\Gamma_2(v)$ *is homogeneous and* $|R_1^*(v^*)| = 2$*, then* $R_2^*(v^*) = \emptyset$*.*

Proof. By Fact 2 and above argument, we have $2 \leq |\Gamma_2^*(v^*)| \leq 3$. Suppose on the contrary that $R_2^*(v^*) \neq \emptyset$. Since $diam(G^*) = 2$ and $\Gamma_1(v)$ is homogeneous, every vertex of $R_2^*(v^*)$ is adjacent to every vertex of $R_1^*(v^*)$. In this way all vertices of $R_2^*(v^*)$ are twins, therefore all vertices of $\bigcup_{u^* \in R_2^*(v^*)} u^*$ are twins, and so $|R_2^*(v^*)| = 1$. Let $R_2^*(v^*) = \{r_2^*\}$, $R_1^*(v^*) = \{x^*, y^*\}$, and $\{a^*, b^*\} \subseteq \Gamma_2^*(v^*)$. Therefore, r_2^* is adjacent to the vertices v^*, x^* and y^*. Note that $\Gamma_1(v)$ and $\Gamma_1^*(v^*)$ are cliques, because $\Gamma_1(v)$ is homogeneous and r_2^* is adjacent to x^* and y^*. Thus, r_2^*, x^* and y^* are of type (1K). Since all neighbors of r_2^* and v^* are shared, r_2^* and v^* are adjacent twins. For $r_2^* \neq v^*$, one of them is of type (N). Because r_2^* is of type (1K), v^* is of type (N). We choose arbitrary fixed vertices $v_1, v_2 \in v^*, r_2 \in r_2^*, x \in x^*, y \in y^*, a \in a^*$ and $b \in b^*$ and set $T = \{v_1, a, b\}$. Since $a^* \neq b^*$ and $\Gamma_2(v)$ is homogeneous, one of the vertices a^* and b^*has only one neighbor in $R_1^*(v^*)$ and the other have one or two. Without loss of generality, we can assume that x^* is the only neighbor of a^* in $R_1^*(v^*)$, see Figure 4(b). Thus,

$$r(v_2|T) = (2, 2, 2), \quad r(x|T) = (1, 1, *), \quad r(r_2|T) = (1, 2, 2), \quad r(y|T) = (1, 2, 1),$$

where $*$ is 1 or 2. However, the four above representations are distinct. This yields the contradiction $\beta(G) \leq n - 4$. Therefore, $R_2^*(v^*) = \emptyset$. ■

Lemma 5. *If* $\Gamma_2(v)$ *is homogeneous and* $|\Gamma_2^*(v^*)| = 3$*, then there is exactly one vertex* $a^* \in \Gamma_2^*(v^*)$ *such that* a^* *is adjacent to all vertices of* $R_1^*(v^*)$*.*

Proof. Let $|\Gamma_2^*(v^*)| = 3$. We have seen that $R_1^*(v^*)$ resolves the set $\Gamma_2^*(v^*)$. Suppose on the contrary that there is not any vertex of $\Gamma_2^*(v^*)$ adjacent to all vertices of $R_1^*(v^*)$. Hence, at least one coordinate of the representation of each vertex in $\Gamma_2^*(v^*)$, with respect to $R_1^*(v^*)$ is 2. While every coordinate of $r(v^*|R_1^*(v^*))$ is 1. Therefore, $R_1^*(v^*)$ is a resolving set for $G^*[R_1^*(v^*) \cup \Gamma_2^*(v^*) \cup \{v^*\}]$ with cardinality $n(G^*) - 4$, since $|\Gamma_2^*(v^*) \cup \{v^*\}| = 4$. It follows that $\beta(G^*) \leq n(G^*) - 4$, and Proposition 3 implies that $\beta(G) \leq n - 4$, which is a contradiction. Therefore, there exists a vertex $a^* \in \Gamma_2^*(v^*)$ adjacent to all vertices of $R_1^*(v^*)$. If there exists another vertex $b^* \in \Gamma_2^*(v^*)$ adjacent to all of $R_1^*(v^*)$, then a^* and b^* are twins, since $\Gamma_2^*(v^*)$ is homogeneous. This implies that $a^* = b^*$ while $|\Gamma_2^*(v^*)| = 3$. Therefore, such a vertex in $\Gamma_2^*(v^*)$ is unique. ■

Lemma 6. *If* $\Gamma_2(v)$ *is homogeneous and* $|R_1^*(v^*)| = 2$*, then* $|\Gamma_2^*(v^*)| \leq 2$.

Proof. On the contrary, suppose $|\Gamma_2^*(v^*)| = 3$. By Lemma 5, there exists exactly one vertex $a^* \in \Gamma_2^*(v^*)$ such that a^* is adjacent to all vertices of $R_1^*(v^*)$. Let $R_1^*(v^*) = \{x^*, y^*\}$ and $\Gamma_2^*(v^*) = \{a^*, b^*, c^*\}$. Each one of vertices b^* and c^* has at least one neighbor in $R_1^*(v^*)$ and by Lemma 5, they have exactly one neighbor in $R_1^*(v^*)$. If their neighbors in $R_1^*(v^*)$ are same, then they are twins, since $\Gamma_2^*(v^*)$ is homogeneous. This implies that every pair of vertices $b \in b^*$ and $c \in c^*$ are twins (because $\Gamma_2(v)$ is homogeneous) consequently, $b^* = c^*$, which contradicts $|\Gamma_2^*(v^*)| = 3$. Thus, one of them, say b^*, is (only) adjacent to x^* and the other c^*, is (only) adjacent to y^*, see Figure 4(a) (dotted edges may be exist or not). Now

$$r(v^*|\{b^*, c^*\}) = (2, 2),\ r(x^*|\{b^*, c^*\}) = (1, 2),\ r(y^*|\{b^*, c^*\}) = (2, 1),\ r(a^*|\{b^*, c^*\}) = (*, *),$$

where $*$ is 1 or 2. If $* = 1$, then $r(a^*|\{b^*, c^*\}) = (1, 1)$, and so $V(G^*)\backslash\{a^*, v^*, x^*, y^*\}$ is a resolving set of size $n(G^*)-4$ for G^*, this contradiction yields $* = 2$. Hence $\Gamma_2(v)$ and $\Gamma_2^*(v^*)$ are independent sets, since $\Gamma_2(v)$ is homogeneous.

Since $R_2^*(v^*) = \emptyset$, if v^* is of type (1N), then v^* and a^* are twins and every pair of vertices $v \in v^*$ and $a \in a^*$ are twins (because both a^* and v^* are of type (1N)), and so $v^* = a^*$, that is a contradiction. Therefore, v^* is of type (K). For arbitrary fixed vertices $v_1, v_2 \in v^*, x \in x^*, y \in y^*, a \in a^*, b \in b^*$ and $c \in c^*$ and $T = \{v_1, a, c\}$, we have

$$r(v_2|T) = (1, 2, 2), \quad r(x|T) = (1, 1, 2), \quad r(y|T) = (1, 1, 1), \quad r(b|T) = (2, 2, 2).$$

Hence, $V(G)\backslash\{v_2, x, y, b\}$ is a resolving set for G, with cardinality $n-4$. This contradiction implies that $|\Gamma_2^*(v^*)| \leq 2$. ■

On account of the above results, we need only assume that $|\Gamma_2^*(v^*)| = |R_1^*(v^*)| = 2$ and $|R_2^*(v^*)| \leq 1$.

Proposition 8. *If* $|R_1^*(v^*)| = 2$ *and* $\Gamma_2(v)$ *is homogeneous, then* G^* *satisfies one of the structures* G_4 *to* G_7.

Proof. Let $R_1^*(v^*) = \{x^*, y^*\}$, $\Gamma_2^*(v^*) = \{a^*, b^*\}$. Then G^* is as described in Figure 5. If $a^* \nsim b^*$, then $x^* \sim y^*$ and $x^* \sim b^*$, otherwise $diam(G^*) = 3$, a contradiction. Let G_0 be the path $(a^*, x^*, v^*, y^*, b^*)$. Thus, G^* must be one of the following five graphs:



(a) (b)

Figure 4: $|\Gamma_2^*(v^*)| = 3$ and $|\Gamma_1^*(v^*)| = 3$.

$H_1^* := G_0 + a^*b^*,$

$H_2^* := G_0 + a^*b^* + x^*b^*,$

$H_3^* := G_0 + a^*b^* + x^*y^*,$

$H_4^* := G_0 + a^*b^* + x^*b^* + x^*y^*,$

$H_5^* := G_0 + x^*b^* + x^*y^*.$

We fix the vertices $v \in v^*, x \in x^*, y \in y^*, a \in a^*$ and $b \in b^*$ in each H_i^*, $1 \leq i \leq 5$. Note that $H_1^* \cong C_5$. If $G^* \cong H_1^*$, then all vertices of H_1^* are of (1), otherwise (with a simple computation) $\beta(G) \leq n - 4$. In this case G^* satisfies structure G_4.

When $G^* \cong H_2^*$, x^* and y^* are not of type (K), because $\Gamma_1(v)$ is homogeneous and $x^* \nsim y^*$. Similarly a^* and b^* are not of type (N). If x^* or y^* is of type (N), then $V(G)\backslash\{v, x, y, b\}$ is a resolving set for G, with cardinality $n - 4$. Also v^* of type (N) or (K) yields $V(G)\backslash\{v, x, y, b\}$ or $V(G)\backslash\{v, x, a, b\}$ is a resolving set for G of size $n - 4$, respectively. These contradictions show that G^* satisfies the structure G_5 if $G^* \cong H_2^*$.

Let $G^* \cong H_3^*$. Since $\Gamma_1(v)$ and $\Gamma_2(v)$ are homogeneous, $x^* \sim y^*$ and $a^* \sim b^*$ imply that x^*, y^*, a^*, b^* are not of type (N). If a^* or b^* is of type (K), then $V(G)\backslash\{v, x, y, a\}$ or $V(G)\backslash\{x, y, v, b\}$ is an $(n - 4)$-vertex resolving set for G, respectively. Also v^* of type (N) yields the resolving set, $V(G)\backslash\{x, y, v, b\}$ for G. These contradictions imply that G^* has the structure G_5.

If $G^* \cong H_4^*$ and one of the vertices v^* or y^* is of type (N), then the set $V(G)\backslash\{v, x, y, b\}$ or $V(G)\backslash\{x, y, a, b\}$ is a resolving set for G of cardinality $n - 4$. Thus v^* and y^* are of type (1K). By symmetry, the vertices a^* and b^* are of type (1K), too. If non-adjacent vertices v^* and b^* are of type (K), then the set $V(G)\backslash\{v, x, y, b\}$ is a resolving set of size $n - 4$. Similarly none-adjacent vertices a^* and y^* are not of type (K) simultaneously. Also, if none-adjacent vertices a^* and v^* are of type (K), then $V(G)\backslash\{v, x, y, a\}$ resolves G, which is impossible. Therefore, none-adjacent vertices are not of the same type (K). Moreover, if x^* is of type (N), and y^* or v^* is of type (K), then $V(G)\backslash\{v, x, y, a\}$ is a resolving set for G of cardinality $n - 4$. By the same reason, if x^* is of type (N), then a^* and b^* are not of type (K). Thus, G^* satisfies the structure G_6.

Finally, assume that $G^* \cong H_5^*$. Since $v^* \neq b^*$ and these vertices are none-adjacent twins in H_5^*, at least one of them is of type (K). Hence, v^* is of type (K) and b^* is of type (1N), because $\Gamma_2(v)$ is homogeneous and $a^* \nsim b^*$. If b^* is of type (N), then $V(G)\backslash\{v,x,y,b\}$ resolves G, a contradiction. It follows that b^* is of type (1). By the similar way as before, one can see that a^* is of type (1), and both x^* and y^* are of type (1K), and thus G^* satisfies the structure G_7. ■



Figure 5: $|\Gamma_1^*(v^*)| = |\Gamma_2^*(v^*)| = 2$ and both $\Gamma_1^*(v^*)$ and $\Gamma_2^*(v^*)$ are homogeneous.

Case 2. For each vertex $v \in V(G)$ with $\Gamma_2^*(v^*) \neq \emptyset$, $\Gamma_1(v)$ is not homogeneous.

We choose a fix vertex $v \in V(G)$ with the property $\Gamma_2^*(v^*) \neq \emptyset$. Lemma 2 concludes that in this case, $\Gamma_2(v)$ is homogeneous. For each vertex $x \in \Gamma_1(v)$, let $M_1(x) := \Gamma_1(v) \cap \Gamma_1(x)$ and $M_2(x) := \Gamma_1(v) \cap \Gamma_2(x)$. Since $M_2(x) \subseteq \Gamma_2(x)$ and $\Gamma_2(x)$ is homogeneous, $M_2(x)$ is also homogeneous. If $M_1(x)$ is not homogeneous, then there exist vertices i, j, and k in $M_1(x)$ such that $i \sim j$ and $k \nsim j$. Thus, for each pair of vertices $y \in M_2(x)$ and $c \in \Gamma_2(v)$, we have $r(i|\{v,x,j\}) = (1,1,1)$, $r(k|\{v,x,j\}) = (1,1,2)$, $r(y|\{v,x,j\}) = (1,2,*)$, $r(c|\{v,x,j\}) = (2,*_1,*_2)$, where $*$, $*_1$ and $*_2$ are 1 or 2. However, these representations are distinct, which is a contradiction. Therefore, $M_1(x)$ is homogeneous.

Proposition 9. *If there exist a vertex $x \in R_2(v)$ with $M_2(x) \neq \emptyset$, then G^* satisfies the structure G_6.*

Proof. Since $x \in R_2(v)$, we have $\Gamma_1(x) = M_1(x) \cup \{v\}$. Note that v is adjacent to all vertices of $M_1(x)$. Since $M_1(x)$ is homogeneous and $\Gamma_1(x)$ is not homogeneous, we conclude $M_1(x)$ is an independent set and contains at least two vertices. Now let m_1 and m_2 be two arbitrary vertices in $M_1(x)$. Thus, m_1 resolves m_2 and v, hence m_1 can not resolve any pair of vertices in $\Gamma_2(x)$, otherwise the set $\{x, m_1\}$ resolves at least four vertices. Therefore, m_1 is adjacent to all vertices of $\Gamma_2(x)$ or none-adjacent to all of them. Since m_1 is an arbitrary vertex of $M_1(x)$, all vertices of $\Gamma_2(x)$ have same neighbors in $M_1(x)$. Note that $\Gamma_2(x) = M_2(x) \cup \Gamma_2(v)$, because $x \in R_2(v)$. Also all vertices of $M_2(x)$ are adjacent to v, and all vertices of $\Gamma_2(v)$ are not adjacent to v. Thus, every pair of vertices in $M_2(x)$ and also every pair of vertices of $\Gamma_2(v)$ are twins. Let $t^* = M_2(x)$ and $s^* = \Gamma_2(v)$ be the corresponding vertices in G^*. Moreover, the vertices of $M_1(x)$ that are adjacent to all of $\Gamma_2(x)$ are twins and form a vertex y^* in G^*, also the remaining vertices of $M_1(x)$ are twins with each other and create a vertex z^* in G^*. Therefore, G^* has at most six vertices v^*, x^*, y^*, z^*, t^*, and s^*, where x^* is adjacent to v^*, z^*, and y^*. Also, v^* and y^* are adjacent to all vertices except s^* and z^*, respectively. There is no other edge in G^* except possibly s^*t^*, see Figure 6(a).

If all of these six vertices exist, then $d(z^*, s^*) = 3$, which contradicts $diam(G^*) = 2$. Since $s^* = \Gamma_2(v)$, the vertex z^* does not exist. It is clear that y^* is of type (N), because $M_1(x)$ is an independent set of size at least two. Let $v \in v^*$, $x \in x^*$, $y_1, y_2 \in y^*$, $s \in s^*$, and $t \in t^*$. If $s^* \nsim t^*$, then $v^* \cup x^* \cup t^* \subseteq \Gamma_2(s)$. But the set $v^* \cup x^* \cup t^*$ is not homogeneous, and so $\Gamma_2(s)$ is not homogeneous, this contradiction yields $s^* \sim t^*$. Thus, since $s^* \cup t^* \subseteq \Gamma_2(x)$, the adjacent vertices s^* and t^* are of type (1K). Moreover, $x^* \cup v^* \subseteq \Gamma_2(s)$ and $x^* \sim v^*$ yields x^* and v^* are of type (1K). But as we see before, when y^* is of type (N) the other vertices can not be of type (K), hence they must be of type (1), also, two none-adjacent vertices are not of the same type (K). Therefore, G^* satisfies the structure G_6. ■

Now let for each vertex $u \in R_2(v)$, one of the sets $M_1(u)$ or $M_2(u)$ is empty. Note that every vertex of $R_2(v)$ has a neighbor in $R_1(v)$, otherwise $diam(G) \geq 3$. Hence, $M_1(u) \neq \emptyset$. Consequently, $M_2(u) = \emptyset$, for each $u \in R_2(v)$. Since $diam(G) = 2$, $M_1(u) = \Gamma_1(v) \backslash \{u\}$. Therefore, every vertex of $R_2(v)$ is adjacent to all vertices of $R_1(v)$, $R_2(v)$ is a clique, and $|R_2^*(v^*)| \leq 1$. We consider the cases $R_1(v)$ is homogeneous or not homogeneous, separately.

Figure 6: $|\Gamma_1^*(v^*)| = 4$ and $|\Gamma_1^*(v^*)| = 2$ in Case 2.

Proposition 10. *If for each vertex $u \in R_2(v)$, the set $M_2(u) = \emptyset$ and $R_1(v)$ is homogeneous, then G^* satisfies the structure G_2.*

Proof. Let $H = G[\{v\} \cup R_1(v) \cup \Gamma_2(v)]$. It is clear that H is an induced subgraph of G with diameter two. By Corollary 1, $n(H) - 3 \leq \beta(H) \leq n(H) - 2$. First suppose $\beta(H) = n(H) - 2$. Theorem A yields H is one of the graphs $K_{r,s}$, $K_r \vee \overline{K}_s$, or $K_r \vee (K_1 \cup K_s)$. If $H = K_{r,s}$, then $R_1(v)$ and $\Gamma_2(v)$ are independent sets. Now, for each $t \in \Gamma_2(v)$ the set $\Gamma_1(t)$ is a nonempty independent set in G, which is a contradiction with the assumption that $\Gamma_1(t)$ is not homogeneous, for such a t. Consequently, $H \neq K_{r,s}$. Note that $R_2(v)$ is a clique and all vertices of $R_2(v)$ are adjacent to all vertices of $R_1(v)$, since for each vertex $u \in R_2(v)$ the set $M_2(u)$ is empty. Therefore, $R_1(v)$ is not a clique, otherwise $\Gamma_1(v)$ is homogeneous. On the other way, if $H = K_r \vee \overline{K}_s$ or $H = K_r \vee (K_1 \cup K_s)$, then $R_1(v)$ is a clique. This contradiction yields $\beta(H) = n(H) - 3$.

Since $R_1(v)$ and $\Gamma_2(v)$ are homogeneous, the graph H with its vertex v satisfies the conditions of Case 1. Thus, H^* has one of the six structures G_2 to G_7. If $R_2(v) = \emptyset$, then $\Gamma_1(v) = R_1(v)$ is homogeneous, which is a contradiction. Therefore, $R_2(v) \neq \emptyset$. Hence, we have $R_2(v)$ is a clique and all its vertices are adjacent to all vertices of $R_1(v)$, consequently, $R_1^*(v^*)$ is not a clique, otherwise $\Gamma_1(v)$ is homogeneous. Note that, in the structures G_3, G_6, and G_7, $R_1^*(v^*)$ is a clique, therefore these structures do not occur.

In graphs with structure G_5, when both neighbors of v^* have degree 3, $R_1^*(v^*)$ is a clique. This implies that, if H^* has the structure G_5, then v^* is adjacent to a degree-2 vertex and a degree-3 vertex. Here, if $t^* \in \Gamma_2^*(v^*)$ is the degree-2 vertex and $t \in t^*$, then $\Gamma_1(t)$ is a clique, contradicts the assumption that $\Gamma_1(v)$ is not homogeneous. Therefore, H^* has not the structure G_5.

When H^* has the structure G_4, $H^* \cong C_5$. If a^* is a neighbor of v^*, then $R_2(v) \cup a^*$ is a resolving set for G with cardinality $n-4$. These contradictions imply that H^* can only have the structure G_2.

Let H^* satisfies the structure G_2. If H^* has not the condition (a) of the structure G_2, then the only vertex of $R_1^*(v^*)$ is of type (1K), therefore, $R_1(v)$ is a clique or $|R_1(v)| = 1$, hence $\Gamma_1(v)$ is a clique, which is impossible. Thus, H^* has the condition (a) of the structure G_2, consequently, the degree-2 vertex of H^* is of type (N), hence, $R_1(v)$ is an independent set of size at least 2. If v^* as a vertex of H^* is of type (N), then by condition (a), the other leaf of H^* is of type (K) and consequently, $V(G)\backslash\{v, r_1, r_2, b\}$ resolves G, where $v \in v^*$, $r_1 \in R_1(v)$, $r_2 \in R_2(v)$, and $b \in \Gamma_2(v)$. Therefore, v^* as a vertex of H^* is of type (1K). Hence, since all vertices of $R_2(v)$ are adjacent to all vertices of $\Gamma_1(v) \cup \{v\}$, v is twin with all vertices of $R_2(v)$. Consequently, v^* as a vertex of G^* is of type (1K) and G^* satisfies the structure G_2. ■

We investigate the case $R_1(v)$ is not homogeneous for two possibility, $|\Gamma_2^*(v^*)| \geq 2$ and $|\Gamma_2^*(v^*)| = 1$, separately.

Proposition 11. *If for each $u \in R_2(v)$ the set $M_2(u) = \emptyset$, $R_1(v)$ is not homogeneous, and $|\Gamma_2^*(v^*)| \geq 2$, then G^* satisfies the structure G_6.*

Proof. Since $\Gamma_2(v)$ is homogeneous, $V(G^*)\backslash\Gamma_2^*(v^*)$ is a resolving set for G^*. Hence, Proposition 3 implies that $\Gamma_2^*(v^*)$ has at most three vertices. Note that, there exist a vertex $x \in R_1(v)$ such that both sets $M_1(x) \cap R_1(v)$ and $M_2(x) \cap R_1(v)$ are nonempty sets, because $R_1(v)$ is not homogeneous. We have $M_1(x) \cap R_1(v)$ and $M_2(x) \cap R_1(v)$ are homogeneous, because $M_1(x)$ and $M_2(x)$ are homogeneous. On the other hand, all vertices of $R_2(v)$ are adjacent to all vertices of $R_1(v)$, since for each vertex $u \in R_2(v)$ the set $M_2(u)$ is empty. Also, $(M_2^*(x) \cap R_1^*(v^*)) \cup \Gamma_2^*(v^*)$ resolves $M_1^*(x) \cap R_1^*(v^*)$, because all different neighbors of the set $M_1^*(x) \cap R_1^*(v^*)$ are in the set $(M_2^*(x) \cap R_1^*(v^*)) \cup \Gamma_2^*(v^*)$. Moreover, in the representations of all vertices of $M_1^*(x) \cap R_1^*(v^*)$ with respect to $\Gamma_2^*(v^*)$, at least one of the coordinates is 1. While all coordinates of $r(v^*|\Gamma_2^*(v^*)$ are 2. Hence, $(M_2^*(x) \cap R_1^*(v^*)) \cup \Gamma_2^*(v^*)$ resolves $\{v^*\} \cup (M_1^*(x) \cap R_1^*(v^*))$. Now Proposition 3 gives $|\{v^*\} \cup (M_1^*(x) \cap R_1^*(v^*))| \leq 3$, and consequently $|M_1^*(x) \cap R_1^*(v^*)| \leq 2$. Similarly $|M_2^*(x) \cap R_1^*(v^*)| \leq 2$.

Since all neighbors of $\Gamma_2^*(v^*)$ are in $R_1^*(v^*)$ and $\Gamma_2(v)$ is homogeneous, $|\Gamma_2^*(v^*)| \geq 2$ implies that there exist vertices $z^* \in R_1^*(v^*)$ and $t^* \in \Gamma_2^*(v^*)$ such that $z^* \nsim t^*$. Let $z \in z^*$. Then $z \nsim t$, for each $t \in t^*$. Since z has a neighbor $t' \in \Gamma_2(v)$, z is adjacent to all vertices of $R_1(v)\backslash\{z\}$ or not adjacent to all these vertices, otherwise the set $\{v, z\}$ resolves four vertices of G. Moreover, if $R_1(v)\backslash\{z\}$ is not homogeneous, then there exist three vertices $i, j, k \in R_1(v)\backslash\{z\}$ such that j resolves $\{i, k\}$, and so $\{v, z, j\}$ resolves $\{i, k, t, t'\}$, which is impossible. Thus, $R_1(v)\backslash\{z\}$ is homogeneous. Therefore, $G[R_1(v)] \cong K_1 \vee \overline{K_l}$ or $K_1 \cup K_l$ for some positive integer $l \geq 2$, because $R_1(v)$ is not homogeneous. It follows that all vertices of $R_1(v)\backslash\{z\}$ have a neighbor and a none-neighbor vertex in $R_1(v)$. Hence, each vertex of $R_1(v)\backslash\{z\}$ is adjacent or none-adjacent to all vertices of $\Gamma_2(v)$, since $\beta(G) = n-3$. But by definition of $R_1(v)$, each vertex of $R_1(v)$ has a neighbor in $\Gamma_2(v)$. From this, each vertex of $R_1(v)\backslash\{z\}$ is adjacent to all vertices of $\Gamma_2(v)$. Thus, all vertices of $R_1(v)\backslash\{z\}$ are twins, and consequently they form a vertex y^* of type (KN) in G^*, and z^* is a vertex of type (1) in G^*. Therefore, $|R_1^*(v^*)| = 2$ and y^* is adjacent to all vertices of $\Gamma_2^*(v^*)$.

If $|\Gamma_2^*(v^*)| = 3$, then z^* can adjacent to one or two vertices of $\Gamma_2^*(v^*)$, however two vertices of $\Gamma_2^*(v^*)$ coincide, because $\Gamma_2(v)$ is homogeneous. Hence, $|\Gamma_2^*(v^*)| = 2$. Let $\Gamma_2^*(v^*) = \{r^*, s^*\}$. Then $G^*[V(G^*)\backslash R_2^*(v^*)]$ is as illustrated in Figure 6(b). If both edges y^*z^* and r^*s^* do not exist, then $d(r^*, z^*) = 3$, which contradicts $diam(G^*) = 2$, therefore one of them exists. Let $y \in y^*, r \in r^*$,-$s \in s^*$. If $y^* \sim z^*$ and $r^* \nsim s^*$, then y^* is of type (N), since $\Gamma_1(v)$ is not homogeneous. Also, r^* is of type (K), otherwise $\Gamma_1(r) = y^*$ is an independent set, which is impossible. This this leads to the $(n-4)$-vertex resolving set, $V(G)\backslash\{v, y, z, r\}$, for G. This contradiction shows that $r^* \sim s^*$ in G^*.

If $y^* \nsim z^*$, then y^* is of type (K), since $\Gamma_1(v)$ is not homogeneous. Moreover, s^* and r^* are of type (1K), since $\Gamma_2(v)$ is homogeneous. However $\Gamma_1(r)$ is a clique, this contradiction shows that both edges r^*s^* and y^*z^* exist in G^*. Therefore, y^* is of type (N), since $\Gamma_1(v)$ is not homogeneous. Furthermore, r^* and s^* are of type (1K), because $\Gamma_2(v)$ is homogeneous. Also, since $\Gamma_2(v)$ is a clique, v^* is of type (1K). As we see before, when y^* is of type (N), the other vertices of G^* are not of type (K), hence other vertices are of type (1). Moreover, two none-adjacent vertices are not of the same type (K). Consequently, G^* satisfies the structure G_6. ■

Proposition 12. *If for each $u \in R_2(v)$ the set $M_2(u) = \emptyset$, $R_1(v)$ is not homogeneous, and $|\Gamma_2^*(v^*)| = 1$, then G^* satisfies one of the structures G_8 to G_{10}.*

Proof. Since $R_1(v)$ is not homogeneous, there exist a vertex $x \in R_1(v)$ such that $M_2(x) \neq \emptyset$. If there is no edge between $R_1^*(v^*) \cap M_1^*(x)$ and $R_1^*(v^*) \cap M_2^*(x)$ or all vertices of $R_1^*(v^*) \cap M_1^*(x)$ are adjacent to all vertices of $R_1^*(v^*) \cap M_2^*(x)$, then $|R_1^*(x)| \leq 3$, because all distinct neighbors of the vertices in these two sets are in each other. In the same manner as the proof of Proposition 11, we can see that each one of the sets $R_1^*(v^*) \cap M_1^*(x)$ and $R_1^*(v^*) \cap M_2^*(x)$ has at most two vertices. Since $R_2(v)$ is a clique and every vertex of it is adjacent to all vertices of $R_1(v)$, if $R_2(v) \neq \emptyset$, then all vertices of $R_2(v)$ are twins with v, and so v^* is of type (K). Also, if $R_2(v) = \emptyset$, then v^* and the only vertex of $\Gamma_2^*(v^*)$, say w^*, are twins. Since $w^* \nsim v^*$, one of them is of type (K). Note that $R_2(v) = \emptyset$ shows v^* is not of type (K), consequently, w^* is of type (K). In this case, by symmetry of v^* and w^* (see Figure 7(a)), we can assume that v^* is of type (K), and so all vertices of $R_2(v)$ are twins with v, and in consequence, without loss of generality we can assume that $R_2^*(v^*) = \emptyset$.

Now let both $R_1^*(v^*) \cap M_1^*(x)$ and $R_1^*(v^*) \cap M_2^*(x)$ have two vertices $\{a^*, b^*\}$ and $\{y^*, z^*\}$, respectively, see Figure 7(a). Since all distinct neighbors of z^* and y^* are in $\{a^*, b^*\}$, $\{a^*, b^*\}$ resolves $\{y^*, z^*\}$. Suppose that $v_1, v_2 \in v^*$, $y \in y^*$, $z \in z^*$, and $w \in w^*$. Then $a^* \cup b^*$ resolves $\{y, z\}$. Also, each coordinate of $r(y|x^*)$ and $r(z|x^*)$ is 2. While $r(v_1|x^*) = r(w|x^*)$ is entirely 1, $r(v_1|v^*\backslash\{v_1\})$ is completely 1, and all coordinates of $r(w|v^*\backslash\{v_1\})$ are 2. Therefore, $V(G)\backslash\{y, z, v_1, w\}$ is a resolving set for G, which is impossible. Hence, at least one of the sets $R_1^*(v^*) \cap M_1^*(x)$ or $R_1^*(v^*) \cap M_2^*(x)$ has one vertex.

If $R_1^*(v^*) \cap M_2^*(x)$ has one vertex and $R_1^*(v^*) \cap M_1^*(x)$ has two vertices, then $a^* \neq b^*$ and $y^* = z^*$. Thus, the only distinct neighbor of a^* and b^* is y^*, and so y^* is adjacent to exactly one of them, say a^*. Now $r(v_1|\{v_2,x,a\})=(1,1,1)$, $r(y|\{v_2,x,a\})=(1,2,1)$, $r(b|\{v_2,x,a\})=(1,1,*)$, and $r(w|\{v_2,x,a\})=(2,1,1)$, where $*=1$ or 2. If $*=2$, then $\beta(G)\leq n-4$, therefore $*=1$. It follows that x^* and b^* are twins and they are adjacent, hence one of them is of type (N). Since $R_1^*(v^*)\cap M_1^*(x)$ is homogeneous, $a^*\sim b^*$ gives b^* is of type (1K), and so x^* is of type (N). In this case, $V(G)\backslash\{x,a,y,w\}$ is a resolving set for G, a contradiction.

If $R_1^*(v^*)\cap M_1^*(x)$ has one vertex and $R_1^*(v^*)\cap M_2^*(x)$ has two vertices, then $a^*=b^*$ and $y^*\neq z^*$. Hence, a^* is adjacent to exactly one of the vertices z^* and y^*, say y^*. Thus, $V(G)\backslash\{v_1,y,z,w\}$ is a resolving set for G. This contradiction yields $|R_1^*(v^*)|\leq 3$. Since $R_1(v)$ is not homogeneous, $|R_1^*(v^*)|\geq 2$. First, let $|R_1^*(v^*)|=3$ and $R_1^*(v^*)=\{x^*,a^*,y^*\}$. Then G^* is as Figure 7(b). If $a^*\nsim y^*$, then x^* and a^* are adjacent twins, consequently, one of them, say x^*, is of type (N). This gives the resolving set, $V(G)\backslash\{v_1,x,y,w\}$ for G, therefore $a^*\sim y^*$, and so x^* and y^* are twins, hence, one of them, say x^*, is of type (K). If y^* is of type (KN), then $V(G)\backslash\{a,x,y,w\}$ is a resolving set for G, thus y^* is of type (1). By the same argument w^* is of type (1). Note that a^* is of type (1K), otherwise $V(G)\backslash\{v_1,y,a,w\}$ is a resolving set for G. Thus G^* satisfies the structure of G_{10}.

When $|R_1^*(v^*)|=2$, two cases can be happened.

1. All vertices of $R_1(v)\cap M_1(x)$ are twins with x. In this case $G^*\cong C_4$ and $x^*=a^*$. Since $R_1(v)\cap M_1(x)$ is not empty, x^* is of type (K). If y^* is of type (KN), then $V(G)\backslash\{v_1,x,y,w\}$ is a resolving set for G, hence y^* is of type (1). By the same reason w^* is of type (1), consequently G^* satisfies the structure G_9.

2. All vertices of $R_1(v)\cap M_2(x)$ are twins with x. In this case G^* is the kite and $x^*=y^*$. Since $R_1(v)\cap M_2(x)$ is not empty, x^* is of type (N). If a^* is of type (N), then $V(G)\backslash\{v_1,x,a,w\}$ is a resolving set for G, hence a^* is of type (1K). If w^* is of type (KN), then $V(G)\backslash\{v_1,x,a,w\}$ is a resolving set for G. Thus, G^* satisfies the structure G_8, which proves the proposition. ■

Now, the proof of necessity is completed.

Figure 7: $|\Gamma_2^*(v^*)|=1$ and $R_1(v)$ is not homogeneous.

3.2 Proof of Sufficiency

In this section we prove that, if G is a graph of order n and $diam(G)=2$ such that G^* has one of the structures G_1 to G_{10} in Theorem 1, then $\beta(G)=n-3$. In the following we consider each structure G_1 to G_{10}, as shown in Figure 1, separately. In each case, we assume that $i \in i^*$, $j \in j^*$, $c \in c^*$, $p \in p^*$, $q \in q^*$, and $u \in u^*$.

$\mathbf{G_1}$. Since G^* has three vertices, by Proposition 4, $\beta(G) \geq n-3$. On the other hand, by Theorem A, $\beta(G) \neq n-2$, also G is not a complete graph. Therefore, $\beta(G) \leq n-3$. Hence $\beta(G) = n-3$.

$\mathbf{G_2}$. Similar to above, we deduce $\beta(G) = n-3$.

$\mathbf{G_3}$. Let H, H_1, H_2, and H_3 be four graphs such that their twin graphs are as G_3 in Figure 1. Also assume that in H^* the vertex j^* is of type (N) and the other vertices are of type (1), in H_1^* both i^* and u^* are of type (K), j^* is of type (N), and p^* is of type (1), in H_2^* both j^* and p^* are of type (N), i^* is of type (K), and u^* is of type (1), in H_3^*, u^* is of type (1) and other vertices are of type (N).

Thus, H is an induced subgraph of G and G is an induced subgraph of H_t, for some t, $1 \leq t \leq 3$. Now we get the metric dimension of H and H_t's, for $1 \leq t \leq 3$. Since in H^*, the vertex j^* is of type (N), each resolving set for H contains at least $|j^*| - 1$ vertices of j^*. Moreover, $r(u|j^*\backslash\{j\}) = r(i|j^*\backslash\{j\})$, hence, $j^*\backslash\{j\}$ is not a resolving set for H. It is easy to check that $(j^*\backslash\{j\}) \cup \{u\}$ is a resolving set, and so a basis of H. Thus, $\beta(H) = n(H) - 3$. Since in H_1^* the vertices i^*, j^*, and u^* are not of type (1), each resolving set for H_1 contains at least $|i^*| - 1, |j^*| - 1$, and $|u^*| - 1$ vertices of i^*, j^*, and u^*, respectively. On the other hand $r(u|i^* \cup j^* \cup u^*\backslash\{i,j,u\}) = r(i|i^* \cup j^* \cup u^*\backslash\{i,j,u\})$, therefore $i^* \cup j^* \cup u^*\backslash\{i,j,u\}$ does not resolve H_1. It is easy to see that $i^* \cup j^* \cup u^*\backslash\{j,u\}$ resolves H_1, hence, it is a basis of H_1, and consequently $\beta(H_1) = n(H_1) - 3$. By a same argument, we can see $\beta(H_t) = n(H_t) - 3$, $2 \leq t \leq 3$. Now, since $diam(H) = diam(G) = 2$, by Corollary 2, we have $\beta(G) = n-3$.

$\mathbf{G_4}$. It is clear that $\beta(C_5) = 2$.

$\mathbf{G_5}$. Let H and R be two graphs such that their twin graphs are as G_5, all vertices of H^* are of type (1) and in R^* both vertices p^* and q^* are of type (1) and other vertices are of type (K). Therefore, H is an induced subgraph of G and G is an induced subgraph of R. It is clear that $\beta(H) = 2 = n(H)-3$. Each resolving set for R contains at least $|i^*|-1$, $|j^*|-1$, and $|u^*|-1$ vertices from i^*, j^*, and u^*, respectively. Let $W = i^* \cup j^* \cup u^*\backslash\{i,j,u\}$. Then $r(i|W) = r(j|W) = r(u|W)$, hence, W is not a resolving set for R. Also adding one of the vertices i, j, and u to W, can not provide a resolving set for R, because $\{i,j,u\}$ is a clique in R. Since $diam(R) = 2$, neither p nor q can not resolve more than two vertices from the set $\{i,j,u\}$. Thus, $\beta(R) \geq |W| + 2 = n(R) - 3$. Since R is not complete graph and none of the graphs in Theorem A, $\beta(R) \leq n(R) - 3$. Hence, $\beta(R) = n(R) - 3$. Since $diam(H) = diam(G) = 2$, Corollary 2 yields $\beta(G) = n-3$.

$\mathbf{G_6}$. Let H, H_1, H_2, and H_3 be four graphs with twin graphs as G_6 in Figure 1. Moreover, assume that, all vertices of H^* are of type (1), in H_1^* the vertex i^* is of type (N) and the other vertices are of type (1), in H_2^* both p^* and q^* are of type (1) and the other vertices are of type (K), in H_3^* both p^* and u^* are of type (1) and the other vertices are of type (K). Hence, H is an induced subgraph of G and G is an induced subgraph of H_t, for some t, $1 \leq t \leq 3$. It is clear that $\beta(H) = 2 = n(H) - 3$. Each resolving set for H_1 contains at least $|i^*| - 1$ vertices from i^*. But

$r(u|i^*\backslash\{i\}) = r(j|i^*\backslash\{i\}) = r(p|i^*\backslash\{i\}) = r(q|i^*\backslash\{i\})$ and there is no vertex of set $\{u, j, p, q\}$ such that adding it to $i^*\backslash\{i\}$ provides a resolving set for H_1, hence $\beta(H_1) \geq |i^*| + 1 = n(H_1) - 3$. Since H_1 is not complete graph, Theorem A gives $\beta(H_1) \leq n(H_1) - 3$, and so $\beta(H_1) = n(H_1) - 3$. Each resolving set for H_2 contains at least $|i^*| - 1, |j^*| - 1$, and $|u^*| - 1$ vertices from i^*, j^*, and u^*, respectively. Assume $W = i^* \cup j^* \cup u^*\backslash\{i, j, u\}$. It follows that $r(i|W) = r(j|W) = r(u|W)$, hence W does not resolve H_2. It is easy to check that to provide a resolving set for H_2 we need to add at least two vertices from $V(H_2) - W$ to W. Thus, $\beta(H_2) \geq |W| + 2 = n(H_2) - 3$. On the other hand, Theorem A implies $\beta(H_2) \leq n(H_2) - 3$, and consequently $\beta(H_2) = n(H_2) - 3$. By the same way, $\beta(H_3) = n(H_3) - 3$. Since $diam(H) = diam(G) = 2$, Corollary 2 implies $\beta(G) = n - 3$.

$\mathbf{G_7}$. Let H and R be two graphs such that their twin graphs are as G_7 in Figure 1. Moreover, assume that in H^* the vertex u^* is of type (K) and the other vertices are of type (1), and in R^* both vertices p^* and q^* are of type (1) and other vertices are of type (K). Therefore, H is an induced subgraph of G and G is an induced subgraph of R. It is clear that each resolving set for H contains at least $|u^*| - 1$ vertices of u^*. Moreover, $r(u|u^*\backslash\{u\}) = r(i|u^*\backslash\{u\}) = r(j|u^*\backslash\{u\})$ and to provide a resolving set for H, we must add at least two vertices from the set $\{u, i, j, p, q\}$ to $u^*\backslash\{u\}$, hence $\beta(H) \geq |u^*| + 1 = n(H) - 3$. Also by Theorem A, $\beta(H) \leq n(H) - 3$, thus $\beta(H) = n(H) - 3$. Since i^*, j^*, and u^* are not of type (1) in R^*, each resolving set for R contains at least $|i^*| - 1$, $|j^*| - 1$, and $|u^*| - 1$ vertices of i^*, j^*, and u^*, respectively. For $W = i^* \cup j^* \cup u^*\backslash\{i, j, u\}$, we have $r(i|W) = r(j|W) = r(u|W)$, hence W is not a resolving set for R. To provide a resolving set for R, we need to add at least two vertices from the set $\{u, i, j, p, q\}$ to W, and consequently, $\beta(R) \geq |W| + 2 = n(R) - 3$. Theorem A shows that $\beta(R) \leq n(R) - 3$, hence $\beta(R) = n(R) - 3$. Since $diam(H) = diam(G) = 2$, Corollary 2 yields $\beta(G) = n - 3$.

$\mathbf{G_8}$. Let H and R be two graphs such that their twin graphs are as G_8 in Figure 1, where in H^* both j^* and p^* are of type (1), u^* is of type (K) and i^* is of type (N), and in R^* both vertices j^* and u^* are of type (K), i^* is of type (N) and p^* is of type (1). Then G is one of the graphs H or R. Theorem A shows that $\beta(H) \leq n(H) - 3$ and $\beta(R) \leq n(R) - 3$. Note that each resolving set for H contains at least $|u^*| - 1$ and $|i^*| - 1$ vertices from u^* and i^*, respectively. If $S = (i^* \cup u^*)\backslash\{i, u\}$, then $r(u|S) = r(j|S)$. Therefore, $\beta(H) \geq |S| + 1 = n(H) - 3$, and so $\beta(H) = n(H) - 3$. It is clear that every resolving set for R contains at least $|i^*| - 1$, $|j^*| - 1$, and $|u^*| - 1$ vertices of i^*, j^*, and u^*, respectively. Let $W = i^* \cup j^* \cup u^*\backslash\{i, j, u\}$. Then $r(j|W) = r(u|W)$, hence W is not a resolving set for R, and so $\beta(R) \geq |W| + 1 = n(R) - 3$. It follows that $\beta(R) = n(R) - 3$. Consequently, $\beta(G) = n(G) - 3$.

$\mathbf{G_9}$. Let G^* be as G_9 in Figure 1, where i^* and u^* are of type (K) in G^*, and the other vertices are of type (1). Each resolving set for G contains at least $|i^*| - 1$ and $|u^*| - 1$ vertices of i^* and u^*, respectively. For $W = i^* \cup u^*\backslash\{i, u\}$, we have $r(i|W) = r(u|W)$, hence W is not a resolving set for G, and so $\beta(G) \geq |W| + 1 = n(G) - 3$. Theorem A implies $\beta(G) \leq n(G) - 3$. Consequently, $\beta(G) = n(G) - 3$.

$\mathbf{G_{10}}$. Let H and R be two graphs such that their twin graphs are as G_{10} in Figure 1, furthermore, in H^* both u^* and c^* are of type (K) and other vertices are of type (1), and in R^* both vertices i^* and p^* are of type (1) and other vertices are of type (K). Then G is one of the graphs H or R.

Theorem A shows that $\beta(H) \leq n(H)-3$ and $\beta(R) \leq n(R)-3$. Note that each resolving set for H contains at least $|u^*|-1$ and $|c^*|-1$ vertices from u^* and c^*, respectively. If $S = c^* \cup u^* \backslash \{c, u\}$, then $r(u|S) = r(j|S) = r(c|S)$, therefore S does not resolve H. To provide a resolving set for H, we need to add at least two vertices from the set $\{u, i, j, c, p\}$ to S, and so $\beta(H) \geq |S|+2 = n(H)-3$, hence $\beta(H) = n(H)-3$. It is clear that every resolving set for R contains at least $|u^*|-1$,- $|j^*|-1$, and $|c^*|-1$ vertices from u^*, j^*, and c^*, respectively. Let $W = u^* \cup j^* \cup c^* \backslash \{u, j, c\}$. Hence, $r(u|W) = r(j|W) = r(c|W)$, therefore W is not a resolving set for R. Clearly, to provide a resolving set for R, we must add at least two vertices from the set $\{u, i, j, c, p\}$ to W, hence $\beta(R) \geq |W|+2 = n(R)-3$, thus $\beta(R) = n(R)-3$. Consequently, $\beta(G) = n(G)-3$.
This completes the proof of the sufficiency of Theorem 1. ■

References

[1] *R.F. Bailey, and P.J. Cameron, Base size, metric dimension and other invariants of groups and graphs,* Bulletin of the London Mathematical Society (to appear).

[2] *R.C. Brigham, G. Chartrand, R.D. Dutton, and P. Zhang, On the dimension of trees,* Discrete Mathematics **294** (2005) 279-283.

[3] *J. Caceres, C. Hernando, M. Mora, I. M. Pelayo, M. L. Puertas, and C. Seara, On the metric dimension of some families of graphs,* Electronic Notes in Discrete Mathematics **22** (2005) 129-133.

[4] *J. Caceres, C. Hernando, M. Mora, I.M. Pelayo, M.L. Puertas, C. Seara, and D.R. Wood, On the metric dimension of cartesian products of graphs,* SIAM Journal Discrete Mathematics **21(2)** (2007) 423-441.

[5] *G.G. Chappell, J. Gimbel, and C. Hartman, Bounds on the metric and partition dimensions of a graph,* Ars Combinatorics **88** (2008) 349-366.

[6] *G. Chartrand, L. Eroh, M.A. Johnson, and O.R. Ollerman, Resolvability in graphs and the metric dimension of a graph,* Discrete Applied Mathematics **105** (2000) 99-113.

[7] *G. Chartrand, and P. Zhang, The theory and applications of resolvability in graphs. (A survey),* In Proceeding 34th Southeastern International Conference on Combinatorics, Graph Theory and Computing **160** (2003) 47-68.

[8] *F. Harary, and R.A Melter, On the metric dimension of a graph,* Ars Combinatorics **2** (1976) 191-195.

[9] *C. Hernando, M. Mora, I.M. Pelayo, C. Seara, and D.R. Wood, Extemal graph theory for metric dimension and diameter,* The Electronic Journal of Combinatorics. (2010) #R30.

[10] *S. Khuller, B. Raghavachari, and A. Rosenfeld, Landmarks in graphs,* Discrete Applied Mathematics **70(3)** (1996) 217-229.

[11] *P.J. Slater, Leaves of trees,* Congressus Numerantium **14** (1975) 549-559.

[12] *G. Sudhakara, and A.R. Hemanth Kumar, Graphs with metric dimension two-a characterization,* World Academy of Science, Engineering and Technology **60** (2009) 621-626.

[13] *I. Tomescu, Discrepancies between metric dimension and partition dimension of a connected graph,* Discrete Mathematics **308(22)** (2008) 5026-5031.

[14] *S.V. Yushmanov, Estimate for the metric dimension of a graph in terms of the diameters and the number of vertices,* Vestnik Moskovskogo Universiteta. Seriya I. Matematika, Mekhanika. **103** (1987) 69-70.